


Cervical Spine Trauma



¹ Stronger consideration for CT/MRI imaging should be given towards the following mechanisms of injury: diving, axial load injury, clothes-linings, high risk MVC (roll-over, ejection, death of occupant in vehicle, > 55mph), falls > 10ft or 2-3x pt height, Concern for NAT, traumatic injury above the clavicles, fracture of other levels of the spine. Additionally stronger consideration for CT/MRI imaging should be given towards non-verbal patients or those with developmental delay.

² C-spine can't be cleared in setting of continued pain from distracting injury or alcohol intoxication. Repeat exam 4-6 hours from last analgesic dose in case of distracting injury. If patient with normal C-spine imaging, can be sent to the floor in a collar. Trauma surgery to reassess and clear on floor ³ C-spine may be cleared by ED, PICU, Trauma or Neurosurgery Teams as indicated

C-spine may be cleared by ED, PICO, Trauma or Neurosurgery Teams as indicate


Reviewers

Created by	Department	Creation Date	Version Date	
J. McCarthy, B. Slater, D. Li	PED, Ped Gem Surg	5/2025	5/2025	

https://www.advocatechildrenshospital.com/healthcare-professionals/peds-pathways



Cervical Spine Trauma



¹ Stronger consideration for CT/MRI imaging should be given towards the following mechanisms of injury: diving, axial load injury, clothes-linings, high risk MVC (roll-over, ejection, death of occupant in vehicle, > 55mph), falls > 10ft or 2-3x pt height, Concern for NAT, traumatic injury above the clavicles, fracture of other levels of the spine. Additionally stronger consideration for CT/MRI imaging should be given towards non-verbal patients or those with developmental delay.

² C-spine can't be cleared in setting of continued pain from distracting injury or alcohol intoxication. Repeat exam 4-6 hours from last analgesic dose in case of distracting injury. If patient with normal C-spine imaging, can be sent to the floor in a collar. Trauma surgery to reassess and clear on floor

³ C-spine may be cleared by ED, PICU, Trauma or Neurosurgery Teams as indicated

B ----

References

Reviewers:					
Created by	Department	Creation Date	Version Date		
J. McCarthy, B. Slater, D. Li	PED, Ped Gem Surg	5/2025	5/2025		

https://www.advocatechildrenshospital.com/healthcare-professionals/peds-pathways



Cervical Spine Trauma

Reviewers:

1. **PECARN prediction rule for cervical spine imaging of children presenting to the emergency department with blunt trauma: a multicentre prospective observational study.** Leonard, Julie C et al.The Lancet Child & Adolescent Health, Volume 8, Issue 7, 482 - 490

Created by	Department	Creation Date	Version Date
J. McCarthy, B. Slater, D. Li	PED, Ped Gem Surg	5/2025	5/2025

https://www.advocatechildrenshospital.com/healthcare-professionals/peds-pathways